



No. 9: Protecting Groundwater from Pesticides

Amy E. Brown, Ph.D., Coordinator
Elizabeth Ingianni, M.S., Program Assistant
Pesticide Education and Assessment Programs
Orig. 1989; Last rev. April 2012

BACKGROUND

Once thought to be safe from contamination, groundwater is now a threatened resource. Recharge areas often lie close to the surface and may be significantly affected by agricultural, residential, or industrial activities. Once contaminated, groundwater is difficult or impossible to clean up. Whether you are a commercial operator treating large acreages or a resident treating your home and lawn, you can take the protective measures discussed in this leaflet to prevent contamination of our water resources.

BEFORE APPLICATION

Identify the vulnerability of the site. Pesticide Information Leaflet No. 8 identifies factors that contribute to groundwater vulnerability. Reduced-tillage cropping systems, surface grading, contour planting and strip cropping can reduce pesticide runoff. Dikes or a buffer zone (a border of untreated vegetation) can slow the movement of runoff water and keep

pesticides out of wells, sink-holes, water bodies, and other sensitive areas.

Evaluate the need for a pesticide. Use an Integrated Pest Management (IPM) approach, combining scientifically sound control strategies with monitoring of pest populations. As part of an IPM program, use pesticides only when necessary and only in amounts that will adequately control the pest. Contact your County Extension office for information on IPM programs.

Choose a pesticide that is less likely to leach. Check the label for warnings about potential to leach or to contaminate groundwater.

Evaluate the method and frequency of pesticide application. Using pesticides applied at the lowest concentration that will control the pest and with the fewest number of applications necessary decreases the amount of pesticide and thus the likelihood of groundwater contamination.

Check the well system. Properly seal new wells, and inspect old wells. Check valves to ensure that seals are adequate. This will help keep accidentally contaminated water from entering wells and, eventually, groundwater.

Monitor weather conditions. Unless the label directs that a pesticide should be watered in after application, pesticides should not be applied when a rain event is expected.

DURING APPLICATION

Read and follow label directions. Many pesticide labels list use instructions or precautions designed to avoid groundwater contamination.

Apply the pesticide at the right time. Fewer applications are required if they are carefully timed in relation to the pest's life cycle. Check the label and ask your County Extension educator for help on timing.

Use the correct rate. Do not use more than the labeled rate. Increasing the dosage increases the cost of pest control, contributes to the potential for resistance development by the pest, and increases chances of environmental contamination.

Mix only the amount needed for the job. Leftover pesticide usually cannot be stored or disposed of easily and safely. Mixing only the amount you need will remove the temptation to overapply.

Calibrate and maintain equipment. Regularly calibrating application equipment reduces the chances of over- or under-applying. Check equipment frequently for leaks and malfunctions.

Prevent back-siphoning and spills. Most contamination incidents are thought to have occurred from carelessness at the mixing/

loading site. Be especially careful near wells, streams, storm drains, or other water sources. Mix pesticides at least 50 feet (preferably, 100 feet) away from these water sources. Be sure to follow label directions if a more conservative setback is specified on the product label. Prevent back-siphoning by keeping the end of the hose above the water level in the spray tank. Install a backflow prevention device (air gap or check valve) on the filling pipe. If a spill does occur, use every precaution to protect water sources. Check with your County Extension office for information on handling small spills.

Direct the application properly. Avoid overspraying the target site. This will help to prevent drift and runoff.

Avoid drift. Do not make applications during windy conditions or during temperature inversions. Drift of pesticide off target and over water sources can contribute to contamination.

Avoid application to bare ground. Never apply pesticide to bare ground unless the label allows it. On sites where groundwater is vulnerable, try to choose pesticides that can be applied post-emergence.

Consider using spray additives. Commercially available spray additives may increase retention on the foliage, decreasing the likelihood that pesticide will be removed by rain or irrigation.

Dispose of excess spray according to label directions. Disposing of leftover spray mix by repeatedly spraying it out at the mixing area or other designated site can contribute significantly to contamination. Instead, excess spray mixture must be applied only on a labeled site in accordance with all label directions.

Exercise care with irrigation and chemigation. Special care should be used during chemigation, as the irrigation water may carry pesticides down through the soil into groundwater. Four safeguards can help to avoid contamination: 1) an interlock connecting the chemical injection equipment with the irrigation pump; 2) a check valve between the irrigation pump and the point of injection; 3) a vacuum breaker behind the check valve; and 4) a low-pressure drain behind the check valve.

AFTER APPLICATION

Store pesticides safely. Pesticides should be stored in their original containers in a cool, well-ventilated, locked location away from pumps and water sources.

Dispose of pesticides and containers safely. Triple rinse pesticide containers and return the rinsate to the spray tank. Puncture the container so it cannot be reused, and dispose of it in a licensed sanitary landfill. Follow label directions for proper disposal of leftover pesticides.

Maintain records of pesticide use. Keep records of the date of application, identity and rate of the pesticide applied, weather conditions, and name of the applicator.

Use care with irrigation after pesticide applications. Because many pesticides move downward through the soil with water, it is best to avoid irrigation immediately after pesticide applications unless the label directions require it. Runoff should be avoided by using a minimum amount of irrigation water. Avoiding runoff will reduce soil erosion and pesticide entry into water.

SOURCES

Brown, A.E. (Ed.) 1991. *Maryland Pesticide Applicator Training Series Core Manual*. Cooperative Extension Service, The University of Maryland System, College Park, MD.

Gardner, Ron. *How to Prevent Water Contamination*. Pesticide Environmental Stewardship, Center for Integrated Pest Management. <http://pesticidestewardship.org/water/Pages/PreventContamination.aspx>.

Nesheim, O. M. and F. M. Fishel. *Protecting Water Resources from Agriculture Pesticides*, Publication #PI1, University of Florida IFAS Extension. <http://edis.ifas.ufl.edu/pi001>.

Trautmann, Nancy M. et al. *Pesticides and Groundwater: A Guide for the Pesticide User*. Cornell University Cooperative Extension. <http://psep.cce.cornell.edu/facts-slides-self/facts/pest-gr-gud-grw89.aspx>.

Waldron, Acie C. 1992. *Pesticides and Groundwater Contamination*, Ohio State Extension Bulletin 820. The Ohio State University Extension. <http://ohioline.osu.edu/b820/index.html> (Accessed 17 April 2012).